

Development and Validation of RP-HPLC Method for Simultaneous Determination of Resveratrol and Curcumin in Pure Form

Manish patidar, Department of Pharmaceutical Chemistry, Sunrise University, Bagad Rajput Dist. Alwar 301001 Rajasthan, India
Dr Mrunal Krishnarao Shirsat Department of Pharmaceutical Chemistry, Sunrise University, Bagad Rajput Dist. Alwar 301001 Rajasthan, India, Corresponding author E-mail: manishm.pati123@gmail.com

ABSTRACT

Cancer chemopreventive agents are designed to reduce the incidence of tumorigenesis by intervening at one or more stages of carcinogenesis. Recently, resveratrol, a natural product found in the diet of humans, has been shown to function as a cancer chemopreventive agent. Resveratrol was first shown to act as an antioxidant and antimutagenic agent, thus acting as an anti-initiation agent. Resveratrol belongs to a class of polyphenolic compounds called stilbenes. (3,5,4'-trihydroxy-trans-stilbene) is a natural compound found in red grape skin, Japanese knotweed (*polygonum cuspidatum*), peanuts and blueberries. A naturally occurring polyphenolic phytoalexin Curcumin, [1,7-bis (4-hydroxy-3-methoxyphenyl)-1, 6- heptadiene-3, 5-dione] is the major yellow pigment extracted from turmeric, a commonly used spice, derived from the rhizome of *Curcuma longa* Linn. In animal studies, curcumin has been shown to increase the survival of tumor-bearing rodents by inhibiting tumor growth and impeding metastasis.

At present study a simple, precise, rapid and accurate reverse phase HPLC method developed for the estimation of Resveratrol and Curcumin in combination as well as individually were developed and validated. The retention time of Resveratrol and Curcumin were found to be 2.90 and 4.11 minutes. Detection was carried out at 424 and 306nm. The regression coefficient value of Resveratrol and Curcumin is 0.9904 and 0.9937 which was found to be linear in the detection range. Limit of detection and limit of quantification of Resveratrol was found to be 0.08 µg/ml and 0.32 µg/ml and Curcumin is 0.05 µg/ml and 0.17 µg/ml. Analysis was performed using a C₁₈ column (250 X 4.6 mm) at room temperature in isocratic mode. The mobile phase used was Citric acid (pH adjusted to 3.5): Acetonitrile (40:60) at flow rate of 1.0 ml/min.

Keywords: HPLC, Resveratrol, Curcumin, Validation.

INTRODUCTION

Resveratrol belongs to a class of polyphenolic compounds called stilbenes. Resveratrol (3,5,4'-trihydroxy-trans-stilbene) is a natural compound. It is a powerful antioxidant produced by some plants to protect them against environmental stresses. Resveratrol, a naturally occurring polyphenolic phytoalexin, is present in many plants and fruits, including red grapes, eucalyptus, spruce, blueberries, mulberries, peanuts, and giant knotweed. Also, red wine contains a lot of it. Resveratrol is an effective antioxidant with strong antiinflammatory and antiproliferative properties. Resveratrol is a fat soluble compound that occurs in a trans and a cis configuration.¹ Curcumin is the principal curcuminoid of the popular Indian spice turmeric, which is a member of the ginger family (*Zingiberaceae*). Turmeric's other two curcuminoids are desmethoxycurcumin and bis-desmethoxycurcumin. The curcuminoids are natural phenols that are responsible for the yellow color of turmeric. Curcumin can exist in several tautomeric forms, including a 1,3-diketo form and two equivalent enol forms. The enol form is more energetically stable in the solid phase and in solution.²

Increasing interest in the use of phytochemicals to reduce prostate cancer led to investigate two potential agents, curcumin and resveratrol as cancer chemo preventive agents. Several *In vitro* assays studies using PTEN-CaP8 cancer cells were performed to investigate the combined effects curcumin with resveratrol on (i) cell growth, apoptosis and cell cycle (ii) impact on activated p-Akt, cyclin D1, m-TOR and androgen receptor (AR) proteins involved in tumor progression. And results of these studies indicated positive anticancer effects of these drugs^{3,4}. Thus in the

present investigation simultaneous estimation method for these two drugs in combination was developed and validated. And value was given to develop a simple, sensitive and repeatable estimation method.

EXPERIMENTAL

Preparation of standard stock solution

The stock solution containing 1mg/ml of Curcumin and Resveratrol were prepared in mobile phase. The stock solution store in light resistant containers. Aliquots of Curcumin (5-50 μ g/ml) and Resveratrol (μ g/ml) were prepared in the mobile phase.

Preparation of combine standard stock solution

An accurately weight quantity equivalent to 100mg of Resveratrol and 50mg of Curcumin were transferred to 100 ml volumetric flask the mixture were dissolved with small quantity of mobile phase and diluted upto the mark and sonicated for 10 minutes at room temperature. The final concentration of Stock solution is 1mg/ml. From this stock solution different dilution of Resveratrol-Curcumin between the range 10-100 μ g/ml and 5-50 μ g/ml respectively, concentration solutions were prepared. These prepared mixture of dilutions were used to determine the linearity using peak area calibration. A plot of peak area versus the respective concentration gives the calibration graph.




Fig: 1 structure of Resveratrol




Fig:2 structure of Curcumin

Chromatographic condition

The HPLC system included a Yung lin instrument 900' UV730D solvent delivery system (pump), injector with a 20 μ L loop volume. The class VP 6.01 data station software was utilized for integration. Separation was achieved using a Princeton C18 column (250 X 4.6 mm, 5 μ ID), (Merck, India). The solvent system consists of Citric acid solution (pH 3.5): acetonitrile (40:60 v/v) was pumped isocratically at a flow rate of 1.0 mL/ Min.⁵⁻⁷

RESULTS AND DISCUSSION

The system suitability tests were carried out on freshly prepared standard stock solution of Resveratrol and Curcumin. Parameters that were studied to evaluate the suitability of the system are given in **Table 1**

Table 1: Combination of Resveratrol and Curcumin

Parameter	Resveratrol	Curcumin
Theoretical plates	587.60	1778.5
Tailing factor	1.1818	1.5000
Retention time(min)	2.90	4.11
LOD μ g/ml	0.08	0.05
LOD μ g/ml	0.32	0.17




Fig 3: Representative Chromatogram for Resveratrol and Curcumin combination by HPLC




Fig :4 Calibration curve indicating linearity range of Resveratrol by HPLC




Fig 5: Calibration curve indicating linearity range of Curcumin by HPLC

Chromatograms eluted from the developed method indicated the retention time of Resveratrol and Curcumin was 2.90 and 4.11 min respectively. A mixture of Acetonitrile and Citric acid solution (pH adjust to 3.5) in the ratio of 60:40 was found to be most suitable. The selected mobile phase was efficient in resolving two drugs to obtain well defined peaks free from tailing. In the present developed HPLC method, the sample preparation required less time. A good linear relationship for both the drugs in combination i.e. for Resveratrol ($r^2=0.9904$) was observed

between the concentration in the linear range of 20-100 μ g/ml and that of Curcumin ($r^2=0.9937$) was between linear range of 10-50 μ g/ml.

Analytical Method Validation Parameters

The method was validated as per ICH guidelines (Q2 R1).

Linearity and Range

Linearity of method is its ability, within a given range, to obtain test results that are directly, or through a mathematical transformation, proportional to concentration of analyte. Good linear correlations were obtained between absorbance and concentration in the selected range of Resveratrol and Curcumin 10 –100 and 5-50 μ g/ml.

Precision

The precision of method expresses the degree of scatter between a series of measurements obtained from multiple sampling of the same homogeneous sample under prescribed conditions. Intraday precision refers to the use of analytical procedure within a laboratory over a short period of time using the same operator with the same equipment whereas Interday precision involves estimation of variations in analysis when a method is used within a laboratory on different days, by different analysts. Repeatability (intraday) was assessed by analyzing these three different concentrations of Resveratrol (20, 40, 60 μ g/ml) Curcumin(10,20,30 μ g/ml) three times a day. Intermediate precision (Interday) was established by analyzing these three different concentrations of Resveratrol (20, 40, 60 μ g/ml) Curcumin(10,20,30 μ g/ml), three times a day for at least three different days. The Standard Deviation, % RSD for the intra-day Resveratrol and Curcumin 0.96 and 0.61.

Accuracy

Accuracy of method is the closeness of test results to true value. It was determined by the application of procedure to recovery studies, where known amount of standard is spiked in preanalyzed samples solutions. The % recovery of the Resveratrol and Curcumin 98.37% and 99.15%.

LOD and LOQ

LOD and LOQ were calculated according to the formulae:

For Resveratrol

$$\text{LOD}=3.3 \sigma / S = 0.08 \mu\text{g/ml}$$

$$\text{LOQ}=10 \sigma / S = 0.32 \mu\text{g/ml}$$

For Curcumin

$$\text{LOD}=3.3 \sigma / S = 0.05 \mu\text{g/ml}$$

$$\text{LOQ}=10 \sigma / S = 0.17 \mu\text{g/ml}$$

CONCLUSION

The development method of estimation indicates that it is simple sensitive and repetitive and major advantage being use of inexpensive solvents. The mixture of solvents used indicated well separation of two peaks without any tailing effects. The developed method can be easily applied for routine analysis of these two drugs Resveratrol and Curcumin in free or formulation dosage forms.

ACKNOWLEDGMENTS:

The authors are thankful to Principal GRY institute of pharmacy Borawan for providing all facilities and his kind support to successful completion of Research project.

REFERENCES

1. Afaq F, Adhami VM, Ahmad N. Prevention of short term ultraviolet B radiation- mediated damages by resveratrol in SKH-1 hairless mice. Toxicol Appl Pharmacol 2003;186:28-37.

2. Kolev, Tsonko M.; Velcheva, Evelina A.; Stamboliyska, Bistra A.; Spiteller, Michael (2005). "DFT and experimental studies of the structure and vibrational spectra of curcumin". *International Journal of Quantum Chemistry* 102 (6): 1069–79.
3. Pezzuto JM. Plant-derived anticancer agents. *Biochem Pharmacol* 1997;53:121–33.
4. Narayanan BA, Re GG, Narayanan NK. Gene expression induced by resveratrol in prostate cancer cells: p53 mediated molecular targets. *Int J Cancer* 2003;104:204–12.
5. Krishna Veni Nagappan. et al., 2009 Liquid Chromatography Method for the Simultaneous Determination of Curcumin and Piperine in Food Products Using Diode Array Detection.P.N. 115-118
6. Kiran Sharma a.et al., 2012 Gupta Development and Validation of UV spectrophotometric method for the estimation of Curcumin in Bulk Drug and Pharmaceutical Dosage Forms.P.N. 375-380
7. Gurinder Singh.et al.,2013 Development and validation of a HPLC method for the determination of trans-resveratrol in spiked human plasma.P.N. 130-135
8. Ammon HP, Wahl MA. Pharmacology of Curcuma longa . *Planta Med* 1991;57:1–7
9. Mukhopadhyay A, Bueso-Ramos C, Chatterjee D, Pantazis P, Aggarwal B. Curcumin downregulates cell survival mechanisms in human prostate cancer cell lines. *Oncogene* 2001;20:7597–609
10. Prashanth R et al.2011Estimation of efavirenz in tablet dosage form by RP-HPLC By Research journal of pharmacy volume 04. pn63-6

