

Boron Chemistry in Aqueous Systems and It's Mathematical Modeling

Charanjit Singh, Research Scholar, Department of Mathematics, SunRise University, Alwar, Rajasthan, India
Dr. Brij Pal Singh, Professor, Department of Mathematics, SunRise University, Alwar, Rajasthan, India

Abstract

Boron is an essential trace element widely distributed in natural waters due to geochemical and anthropogenic processes. This paper presents a comprehensive study of boron chemistry in aqueous systems, including its speciation, solubility, environmental behavior, and health effects. Furthermore, a mathematical modeling approach using the least squares method is applied to analyze boron concentration data. A cubic regression model is developed to describe the relationship between observed variables, demonstrating the nonlinear nature of boron distribution in water systems.

Introduction

Boron is a naturally occurring element found in various environmental compartments such as soil, water, and air. It enters aquatic systems primarily through the weathering of boron-containing minerals and anthropogenic activities such as agriculture and industrial discharge. Understanding boron chemistry in water is essential due to its dual role as a micronutrient and a potential toxicant at elevated concentrations.

This study focuses on the chemical behavior of boron in aqueous systems and the development of an effective mathematical model to represent its concentration variations.

Discussion about Boron-values:

Br \$ O2: response contrivances, eco-friendly impression and well-being belongings:

Salt-water comprehends roughly 4.00-5.01 ppm br. 'River-water' largely contains only 10 ppb. In seaweed 8-15 ppm and in mussels 4-5 ppm preliminaries of "br" was found. "Br" melted in water occurs as $B^{-}OH^{-3}$ (aquatic) or $B^{-}OH^{-4}$ (aquatic).

In pardon method and in pardon procedure does br respond with aquatic?

"Br" does not respond with aquatic under normal circumstances. Nevertheless, this may be the event for boron compounds. E.g., the "Br" Trifluoride Ethyl-Ether Multifaceted reacts with aquatic and forms diethyl ether BF_3 . Several boron-compounds are hydrolyzed in water, such as the triodium iodide.

Solubility-of-boron and boron-compounds

In addition, boron-salts are water-soluble. The soluble water content of the boric acids contains 57 g/L, 25.2 g/L boron, and 22 g/L boron trioxide. The least water-soluble boron complex of boron trifluoride is 2.4 g/L water-soluble. Some boron compounds are totally water insoluble, for example, boron nitrite.

Why is boron present in water?

Kernite, borax, ulexite and colemanite are the most common minerals that contain boron. It is also present in rock formations of slate and loam. Boron amounts between 5 and 80 ppm are found in the airtight soil. Rich boron sites include boron, boron, and mineral boron, such as fumaroles. The degree to which tones are connected depends primarily on the pH. Boron is released by weathering from rocks and soils and then ends in water.

Pure elements are never found in manufacturing except in the manufacture of metal borides or to increase the conductivity of aluminium.

The metal borides are manufactured to engines, rocket power, high-temperature reaction containers and electrodes, for example. A boron is applied to the hardness of the material.

Perborate sodium is used in detergents as bleach. This inevitably shapes borate, destroying water plants directly. Borates are used as softening agents for water. The glass, the glass fibre and the ceramics, email and other boron compounds are used. When boron is added and the glass fibres applied as insulation, the glass is thicker and heat resistant. Cleansing chemicals,

batteries, illicit conservators and eye drops can also contain boron compounds. Borax, boron oxide and boric acid are the most common compounds in this regard. In vast quantities of fertilisers and pesticides are added to boric acid and borax. The factor is also present in wood preservatives and impregnation agents. As boron carbide and boron nitrite, it is applied as an abrasive.

In a polymer matrix, boron has a special purpose. It is used as cooling agent in unsafe conditions to control nuclear reactors. In the core of the reactor it also absorbs neutrons.

When Boron is not properly sealed, it also ends in soil and groundwater through domestic landfills. It is a standard compound measure suggesting the presence of other harmful compounds.

What are the environmental effects of boron in water?

For several animals, boron is a dietary necessity and has a significant function to play in mitosis. This is true for green algae and other other species of plants. Boron shortcomings cause problems of growth and complexity in the mobilisation of sugar. The most consumed boron compound is boric acid. Plants have boron 30-75 ppm (dry mass). At concentrations of over 100 ppm the toxic process continues. This will lower the yield of crops. Grass species tolerate moderately high amounts of boron, but pine species are particularly vulnerable. But in contrast with other plant species, trees need significant quantities of boron. About 25 ppm is a tolerable boron in the soils.

High levels of boron in water, for concentrations of 10-300 mg/L, may be harmful to fish species. Mainly borate is unsafe for water plants. The food for vertebrates is not boron.

Boric acid is marginally water-dangering, but it is very water-dangerous for boron halogens.

Boron is mobile on average and is steadily converted. It can therefore spread easily by water.

Two stable and 14 unstable isotopes are made up of the boron.

What are the health effects of boron in water?

The body contains about 0,7 ppm of boron, a nutritional requirement not considered. Nevertheless, we ingest this ingredient from food, as it is plant food necessity. About 2 mg of daily consumption. There is no toxicity level for the volume of boron in fruits and vegetables. For a daily intake of more than 5 g boric acid, nausea, vomiting and diarrhoea and blood clotting are clearly negatively affecting the human body. Menops exceeding 20 g pose a danger to life. The skin and eyes are damaged by boric acid. Boron trifluoride skin contact can lead to corrosion.

The amount of boron in soils and drinking water is possible to associate with the prevalence of arthritis amongst humans.


Boric acid as well as borax are in many concentrations applied in medicine. In brain tumour therapy neutron-absorbing properties of boron are added (boron neutron capture therapy).

Which water purification technologies can be applied to remove boron from water?

Boron is of course, predominantly present in the form of boric acid and boric acid salts. Boric acid, owing to its proximity to silicate, can be taken off by ion exchanger but very slowly.

Table: Boron-Level

Particulars of Elements	Farm water			River water		
	H.P.	Punjab	Rajasthan	H.P.	Punjab	Rajasthan
BORON	0.82	0.77	0.28	0.05	0.14	0.31



Calculate fitting a cubic equation - Curve fitting using least square method:

X	Y
0.82	0.77
0.82	0.28
0.77	0.28
0.05	0.14
0.05	0.31
0.14	0.31

Solution:

The equation is $y=a+bx+cx^2+dx^3$ and the normal equations are

$$\sum y = a\sum 1 + b\sum x + c\sum x^2 + d\sum x^3$$

$$\sum xy = a\sum x + b\sum x^2 + c\sum x^3 + d\sum x^4$$

$$\sum x^2y = a\sum x^2 + b\sum x^3 + c\sum x^4 + d\sum x^5$$

$$\sum x^3y = a\sum x^3 + b\sum x^4 + c\sum x^5 + d\sum x^6$$

The values are calculated using the following table:

x	y	x ²	x ³	x ⁴	x ⁵	x ⁶	x·y	x ² ·y	x ³ ·y
0.82	0.77	0.6724	0.55137	0.45212	0.37074	0.30401	0.6314	0.51775	0.42455
0.82	0.28	0.6724	0.55137	0.45212	0.37074	0.30401	0.2296	0.18827	0.15438
0.77	0.28	0.5929	0.45653	0.35153	0.27068	0.20842	0.2156	0.16601	0.12783
0.05	0.14	0.0025	0.00013	0.00001	0	0	0.007	0.00035	0.00002
0.05	0.31	0.0025	0.00013	0.00001	0	0	0.0155	0.00078	0.00004
0.14	0.31	0.0196	0.00274	0.00038	0.00005	0.00001	0.0434	0.00608	0.00085
$\sum x = 2.65$	$\sum y = 2.09$	$\sum x^2 = 1.9623$	$\sum x^3 = 1.56226$	$\sum x^4 = 1.25617$	$\sum x^5 = 1.01221$	$\sum x^6 = 0.81644$	$\sum x \cdot y = 1.1425$	$\sum x^2 \cdot y = 0.87923$	$\sum x^3 \cdot y = 0.70767$

Substituting these values in the normal equations:

$$6a+2.65b+1.9623c+1.56226d=2.09$$

$$2.65a+1.9623b+1.56226c+1.25617d=1.1425$$

$$1.9623a+1.56226b+1.25617c+1.01221d=0.87923$$

$$1.56226a+1.25617b+1.01221c+0.81644d=0.70767$$

Solving these 4 equations, Total Equations are 4

$$6a+2.65b+1.9623c+1.56226d=2.09 \rightarrow (1)$$

$$2.65a+1.9623b+1.56226c+1.25617d=1.1425 \rightarrow (2)$$

$$1.9623a+1.56226b+1.25617c+1.01221d=0.87923 \rightarrow (3)$$

$$1.56226a+1.25617b+1.01221c+0.81644d=0.70767 \rightarrow (4)$$

Select the equations (1) and (4), and eliminate the variable d.

$$6a+2.65b+1.9623c+1.56226d=2.09 \quad \times 1 \rightarrow \quad 6a \quad +2.65b \quad + \frac{1.9623}{c} \quad + \frac{1.5622}{6d} = 2.09$$

$$1.56226a+1.25617b+1.01221c+0.816 \times 1.913 \quad 2.9893 \quad + \frac{2.4036}{8b} \quad + \frac{1.9368}{7c} \quad + \frac{1.5622}{6d} = \frac{1.354}{13}$$

$$44d=0.70767 \quad 5 \rightarrow$$

$$\frac{3.0106}{1a} \quad + \frac{0.2463}{2b} \quad + \frac{0.0254}{3c} = \frac{0.735}{87} \rightarrow (5)$$

Select the equations (2) and (4), and eliminate the variable d.

$$2.65a+1.9623b+1.56226c+1.25617d \quad \times 1 \rightarrow \quad 2.65a \quad + \frac{1.9623}{b} \quad + \frac{1.562}{26c} \quad + \frac{1.2561}{7d} = \frac{1.142}{5}$$

$$=1.1425$$

$$1.56226a+1.25617b+1.01221c+0.816 \times 1.5385 \quad 2.403 \quad + \frac{1.9327}{4b} \quad + \frac{1.557}{38c} \quad + \frac{1.2561}{7d} = \frac{1.088}{82}$$

$$44d=0.70767 \quad 9 \rightarrow$$

$$\frac{0.246}{32a} \quad + \frac{0.0295}{6b} \quad + \frac{0.004}{88c} = \frac{0.053}{68} \rightarrow (6)$$

Select the equations (3) and (4), and eliminate the variable d.

$$1.9623a+1.56226b+1.25617c+1.0122 \quad \times 1 \rightarrow \quad 1.962 \quad + \frac{1.5622}{6b} \quad + \frac{1.256}{17c} \quad + \frac{1.0122}{1d} = \frac{0.879}{23}$$

$$1d=0.87923$$

$$1.56226a+1.25617b+1.01221c+0.816 \times 1.2397 \quad 1.936 \quad + \frac{1.5573}{8b} \quad + \frac{1.254}{92c} \quad + \frac{1.0122}{1d} = \frac{0.877}{36}$$

$$44d=0.70767 \quad 8 \rightarrow$$

$$\frac{0.025}{43a} \quad + \frac{0.0048}{8b} \quad + \frac{0.001}{25c} = \frac{0.001}{87} \rightarrow (7)$$

Select the equations (5) and (7), and eliminate the variable c.

$$3.01061a+0.24632b+0.02543c=0.7 \quad \times 1 \rightarrow \quad 3.01061 \quad + \frac{0.24632}{b} \quad + \frac{0.02543}{c} = \frac{0.7358}{7}$$

$$3587$$

$$0.02543a+0.00488b+0.00125c=0 \quad \times 20.39083 \quad 0.51861 \quad + \frac{0.09949}{b} \quad + \frac{0.02543}{c} = \frac{0.0381}{6}$$

$$0187 \quad \rightarrow$$

$$2.492a \quad + \frac{0.14682}{b} = \frac{0.6977}{1} \rightarrow (8)$$

Table: Discussion about Boron-values:

The values of EC are in worst position in the farm on the other hand the Boron values of river is null in compare to (4.0-5.0) see (chart-6), and the cubic model as follows:

$$y=0.11041+2.87812x-12.01002x^2+11.11709x^3 - \sigma^2/2 - \delta$$

Discussion

The study reveals that boron concentrations are generally higher in farm water compared to river water due to agricultural inputs. The developed model effectively captures the variation pattern and can be used for environmental assessment and prediction.

Conclusion

Boron is an important element in aquatic systems, exhibiting both beneficial and adverse effects depending on its concentration and chemical form. At low concentrations, boron functions as an essential micronutrient, particularly for plant growth and metabolic activities such as cell division and carbohydrate transport. However, when present in elevated concentrations, it becomes toxic to plants, aquatic organisms, and potentially harmful to human health. This dual role highlights the importance of monitoring and managing boron levels in natural and engineered water systems.

The occurrence of boron in water is governed by a combination of natural and anthropogenic factors. Natural sources include the weathering of boron-containing minerals such as borax, kernite, and ulexite, which release boron into groundwater and surface water. In addition, geological formations and soil composition significantly influence boron mobility and concentration. Anthropogenic activities, including agricultural practices (use of fertilizers and pesticides), industrial discharges, detergent usage, and waste disposal, further contribute to increased boron levels in water bodies. These inputs can lead to localized contamination, particularly in agricultural regions.

From a chemical perspective, boron exists predominantly in aqueous systems as boric acid and borate ions, with their distribution strongly dependent on pH conditions. This speciation plays a crucial role in determining boron's reactivity, mobility, and bioavailability. The relatively high solubility and mobility of boron compounds allow them to spread easily through water systems, making their control and removal a challenging task.

References

1. World Health Organization. (2017). *Guidelines for drinking-water quality* (4th ed.). WHO Press.
2. Hem, J. D. (1985). *Study and interpretation of the chemical characteristics of natural water* (3rd ed.). U.S. Geological Survey.
3. Greenwood, N. N., & Earnshaw, A. (1997). *Chemistry of the elements* (2nd ed.). Butterworth-Heinemann.
4. Cotton, F. A., Wilkinson, G., Murillo, C. A., & Bochmann, M. (1999). *Advanced inorganic chemistry* (6th ed.). Wiley.
5. Weast, R. C. (Ed.). (1984). *CRC handbook of chemistry and physics* (64th ed.). CRC Press.
6. Parks, J. L., Edwards, M., & McNeill, L. S. (2004). Effect of pH, alkalinity, and dissolved solids on boron removal using reverse osmosis. *Journal of Environmental Engineering*, 130(1), 85–92.
7. Nadav, N. (1999). Boron removal from seawater reverse osmosis permeate utilizing selective ion exchange resin. *Desalination*, 124(1–3), 131–135.
8. Kabay, N., Bryjak, M., & Hilal, N. (Eds.). (2010). *Boron separation processes*. Elsevier.
9. Goldberg, S. (1997). Reactions of boron with soils. *Plant and Soil*, 193(1–2), 35–48.
10. Gupta, U. C. (Ed.). (1993). *Boron and its role in crop production*. CRC Press.
11. Camacho-Cristóbal, J. J., Rexach, J., & González-Fontes, A. (2008). Boron in plants: Deficiency and toxicity. *Journal of Integrative Plant Biology*, 50(10), 1247–1255.

12. Eisler, R. (1990). *Boron hazards to fish, wildlife, and invertebrates: A synoptic review*. U.S. Fish and Wildlife Service.
13. Butterwick, L., de Oude, N., & Raymond, K. (1989). Safety assessment of boron in aquatic and terrestrial environments. *Ecotoxicology and Environmental Safety*, 17(3), 339–371.
14. Power, P. P., & Woods, W. G. (1997). The chemistry of boron and its speciation in plants. *Plant and Soil*, 193(1–2), 1–13.
15. U.S. Environmental Protection Agency. (2008). *Health effects support document for boron*. EPA.
16. Kot, F. S. (2009). Boron sources, speciation and its potential impact on health. *Reviews in Environmental Science and Bio/Technology*, 8(1), 3–28.
17. APHA. (2012). *Standard methods for the examination of water and wastewater* (22nd ed.). American Public Health Association.
18. Montgomery, D. C., & Runger, G. C. (2014). *Applied statistics and probability for engineers* (6th ed.). Wiley.

